

File _____

Those Eligible
To Read The
Attached

Copy# 5A
C. Hickman
901

Before reading this document, sign and date below:

[illegible]

Distribution List

1. MTHalley
2. HSMoalcraft
3. Col. McCormick for Lt. HSMoalcraft
4. LEBulet
5. ESWithowski
6. SAReynolds file

This document contains information affecting the National Defense of the United States within the meaning of the espionage laws, the transmission or the revelation of its contents in any manner to an unauthorized person is prohibited and may result in severe civil or criminal penalties under applicable laws.

This document has been approved for release
to the public by:

David R. Hamlin 2/3/95
Technical Information Officer Date
ORNL Site

~~RESTRICTED DATA~~

SECRET

Document consists of _____

Pages and _____

No. _____ of _____

Figures _____

Copies, Series _____

INTER-COMPANY CORRESPONDENCE

~~SECRET~~

OAK RIDGE NATIONAL LABORATORY

Post Office Box P
OAK RIDGE, TENN.

(INSERT
NAME)

COMPANY CARBIDE AND CARBON CHEMICALS CORP. LOCATION

Classification Cancelled

TO
LOCATION

ATTENTION
COPY TO

M. T. ~~Reynolds~~

By Authority Of

DATE

June 9, 1950

By

~~ATG~~

Date

~~AUG 30 1971~~

ANSWERING LETTER DATE

H. J. McAlduff

Col. McCormick for Lt. W. E. Carr

L. E. Emlet

E. J. Witkowski

E. A. Reynolds file

SUBJECT

Gross Beta and Gamma
Activities of Dissolver
Solution.

The gross beta and gamma activities of a typical sample of ORNL dissolver solution (fission iodine process) were determined in order to evaluate the extent of settling basin contamination caused by the Army laundry program in July.

The beta activity was determined by counting a suitable aliquot dried on a one-inch watch glass and supported on the second shelf of the standard counter. The counting efficiency of this arrangement is about 10%. This simulates adequately the conditions of measurement of settling basin beta activity. Decay was followed for two weeks. The figure shows nominal millicuries per milliliter as a function of time after pile discharge.

The gamma activity was measured by means of the gamma ionization chamber. The activity was calculated as 1 Mev photons (see figure).

E. A. Reynolds
E. A. Reynolds

SAR:ko

Classification changed to: Unclassified
(level and category)

By authority of: DAR-1 & PR-2
(classification guide)

WV Bush 4/11/94

ADC or ADD signature (first reviewer) Date

Ted Davis 4/25/94

ADD signature (final reviewer) Date

~~RESTRICTED DATA~~

This document contains restricted data as defined in the Atomic Energy Act of 1946.

~~CAUTION~~

This document contains information affecting the National Defense of the United States. Its transmission or the disclosure of its contents in any manner to an unauthorized person is prohibited and may result in severe criminal penalties under applicable Federal laws.

~~SECRET~~

DECAY OF ACTIVITY OF DISSOLVER SOLUTION

○ BETA
● GAMMA

1 WEEK

2 WEEKS


$\beta \sim 25$

$\beta \sim 16$

$\gamma \sim 10$

$\gamma \sim 7$

DAYS AFTER DISCHARGE



~~SECRET~~

Dup

File _____

Those Eligible
To Read The
Attached

Copy# 32
C. Thakurki
901

Before reading this document, sign and date below:

Name _____ Date _____

8 MAY

54 INT

~~CONFIDENTIAL~~

This document

Publicly Releasable

This document has received the necessary patent and technical information reviews and can be distributed without limitation.

ADD signature 1/18/95
Date

Single rereview of CCRP-declassified documents was authorized by DOE Office of Declassification memo of August 22, 1994.

define

~~SECRET~~

INTER-COMPANY CORRESPONDENCE

OAK RIDGE NATIONAL LABORATORY
 (INSERT NAME) COMPANY CARBIDE AND CARBON CHEMICALS CORP. LOCATION Post Office Box P
 OAK RIDGE, TENN.

TO
 LOCATION

ATTENTION
 COPY TO

M. T. ~~_____~~ DATE June 9, 1950
 By ~~_____~~
 By ATB Date 1/18/95 ANSWERING LETTER DATE

E. J. McAlduff
 Col. McCormick for Lt. W. E. Jarr SUBJECT Gross Beta and Gamma
 L. E. Inlet Activities of Dissolver
 L. J. Witkowski Solution.
 L. A. Reynolds file

The gross beta and gamma activities of a typical sample of CHL dissolver solution (fission iodine process) were determined in order to evaluate the extent of settling basin contamination caused by the Army laundry program in July.

The beta activity was determined by counting a suitable aliquot dried on a one-inch watch glass and supported on the second shelf of the standard counter. The counting efficiency of this arrangement is about 10%. This simulates adequately the conditions of measurement of settling basin beta activity. Decay was followed for two weeks. The figure shows nominal millicuries per milliliter as a function of time after pile discharge.

The gamma activity was measured by means of the gamma ionization chamber. The activity was calculated as 1 Mev photons (see figure).

S. A. Reynolds
 S. A. Reynolds

SAR:ko

CLASSIFICATION CANCELLED

AS Just 1/18/95
 ADD signature Date

Single rereview of CCRP-declassified documents was authorized by DOE Office of Declassification memo of August 22, 1994.

~~RESTRICTED DATA~~
 This document contains restricted data as defined in the Atomic Energy Act of 1946.

~~CAUTION~~

This document contains information affecting the National Defense of the United States. Its transmission or the disclosure of its contents in any manner to an unauthorized person is prohibited by law.

DECAY OF ACTIVITY OF DISSOLVER SOLUTION

